

ON PSEUDO ALGEBRAICALLY CLOSED EXTENSIONS OF FIELDS

LIOR BARY-SOROKER

ABSTRACT. The notion of ‘Pseudo Algebraically Closed (PAC) extensions’ is a generalization of the classical notion of PAC fields. In this work we develop a basic machinery to study PAC extensions. This machinery is based on a generalization of embedding problems to field extensions. The main goal is to prove that the Galois closure of any proper separable algebraic PAC extension is its separable closure. As a result we get a classification of all finite PAC extensions which in turn proves the ‘bottom conjecture’ for finitely generated infinite fields.

The secondary goal of this work is to unify proofs of known results about PAC extensions and to establish new basic properties of PAC extensions, e.g. transitivity of PAC extensions.

1. INTRODUCTION AND RESULTS

This work concerns Pseudo Algebraically Closed (PAC) extensions of fields, and especially the Galois structure of PAC extensions. We start by a short survey on this notion and its importance. Then we discuss the results and methods of this work.

1.1. Pseudo Algebraically Closed Extensions. A field K is called **PAC** if it has the following geometric feature: $V(K) \neq \emptyset$ for any non-void absolutely irreducible variety V which is defined over K . In [15] Jarden and Razon generalize this classical notion to a field K and a subset K_0 (in this work K_0 will always be a field, unless otherwise stated): We call K/K_0 **PAC extension** (or just say that K/K_0 is PAC) if for every absolutely irreducible variety V of dimension $r \geq 1$ which is defined over K and every separable dominating rational map $\nu: V \rightarrow \mathbb{A}^r$ defined over K there exists $a \in V(K)$ for which $\nu(a) \in K_0^r$. (See Proposition 3.1 for equivalent definitions of PAC extensions in terms of polynomials and of places.) For example, every separably closed field is PAC over any infinite subfield, and for every PAC field K , the trivial extension K/K is PAC.

The original motivation of Jarden and Razon for this definition is related to a generalization of Hilbert’s 10th problem to ‘large’ algebraic rings. The problem asks whether there exists an algorithm that determines whether a system of polynomial equations over $\mathbb{Z}$ has a solution in $\mathbb{Z}$. Matijasevich gave a negative answer to that

Date: July 16, 2009.

2000 Mathematics Subject Classification. 12E30, 12F10.

Key words and phrases. Field Arithmetic, PAC, embedding problem.

This work is a part of the author’s PhD thesis done at Tel Aviv University under the supervision of Prof. Dan Haran .

This research was partially supported by the ISRAEL SCIENCE FOUNDATION (Grant No. 343/07).

problem relying on the works of Davis, Putnam, and J. Robinson since the 1930s (see [19]).

A natural generalization of Hilbert's 10th problem is to consider solutions in other rings. In [23] Rumely establishes a local-global principle for the ring $\tilde{\mathbb{Z}}$ of all algebraic integers, and deduces from it a positive answer of Hilbert's 10th problem for that ring. (It is interesting to mention that for the ring $\mathbb{Q}$ the problem is still open.)

In [16] Jarden and Razon extend Rumely's local global principle to the ring of integers R of an algebraic field K , for 'almost all' $K \subseteq \tilde{\mathbb{Q}}$ whose absolute Galois group $\text{Gal}(K)$ is finitely generated. Their key idea is to use the fact (proven by them in [15]) that these fields are PAC over $\mathbb{Z}$, and hence over R . Then, following Rumely, they deduce a positive answer to Hilbert's 10th problem for R .

In [17] Jarden and Razon continue their work, and establish Rumely's local-global principle for smaller rings. They also deal with the positive characteristic case, and strengthen the local-global principle itself.

The applications of PAC extensions are not restricted to Hilbert's 10th problem and Rumely's local-global principle, and indeed some other applications recently appeared in the literature. Some examples are: (1) new constructions of Hilbertian domains [21]; (2) an analog of Dirichlet's theorem on primes in arithmetical progressions for a polynomial ring in one variable over some infinite fields [3]; (3) the study of the question: When is a non-degenerate quadratic form isomorphic to a scaled trace form? [6].

1.2. The Galois Closure of PAC Extensions. In [15] (where PAC extensions first appear) Jarden and Razon find some Galois extensions K of $\mathbb{Q}$ such that K is PAC as a field but K is a PAC extension of no number field. (For this a heavy tool is used, namely Faltings' theorem.) Then they ask whether this is a coincidence or a general phenomenon.

In [14] Jarden settles this question by showing that the only Galois PAC extension of an arbitrary number field is its algebraic closure. Jarden does not use Faltings' theorem, but different results. Namely, Razon's splitting theorem (see Theorem 6), Frobenius' density theorem, Neukirch's characterization of p -adically closed fields among all algebraic extensions of $\mathbb{Q}$, and also the special property of $\mathbb{Q}$ that it has no proper subfields(!). For that reason Jarden's method is restricted to number fields.

The next step is to consider a finitely generated infinite field K_0 . Elaborating the original method of Jarden-Razon, i.e. using Faltings' theorem (and the Grauert-Manin theorem in positive characteristic), Jarden and the author generalize Jarden's result to K_0 [5].

In this work we further generalize this theorem to the most general case, where K_0 is an arbitrary field. Namely we prove that the only Galois PAC extensions are the trivial ones (see Theorem 1 below). Our proof is based on the *lifting property* which will be discussed later in the introduction and the realization of wreath products in fields. Thus it uses no special features of finitely generated fields, and for that reason it applies to any K/K_0 .

Theorem 1. *Let K/K_0 be a proper separable algebraic PAC extension. Then the Galois closure of K/K_0 is the separable closure of K_0 . In particular, if K/K_0 is a Galois PAC extension, then either $K = K_0$ or $K = K_s$.*

It is important to note in this stage, that there are a lot of PAC extensions, which are not Galois. Let e be a positive integer. If K_0 is a countable Hilbertian field, then for almost all $\sigma = (\sigma_1, \dots, \sigma_e) \in \text{Gal}(K_0)^e$ (in the sense of the Haar measure)

$$K_{0s}(\sigma) = \{x \in K_{0s} \mid \sigma_i(x) = x, \forall i\}$$

is a PAC extension of K_0 [15], and hence of any subfield of $K_{0s}(\sigma)/K_0$. Using this result, in [3, 6] many PAC extensions are constructed. For example if K_0 is a pro-solvable extension of a countable Hilbertian field, then there exists a PAC extension K/K_0 such that the order of $\text{Gal}(K)$ (as a supernatural number) is $\prod_p p^\infty$. In a sequel [1], the author studies the group theoretic properties of the pair of profinite groups $\text{Gal}(K) \rightarrow \text{Gal}(K_0)$ and using the transitivity of PAC extensions appear here constructs new PAC extensions. For example, for any projective profinite group P of rank at most countable, $\mathbb{Q}_{ab}$ has a PAC extension with absolute Galois group P . It is open whether a finitely generated field K_0 has a PAC extension whose absolute Galois group is not finitely generated [5], Conjecture 7.

Note that Theorem 1 generalizes the following

Theorem 2 (Chatzidakis [10], Theorem 24.53). *Let K_0 be a countable Hilbertian field. Then for almost all $\sigma \in \text{Gal}(K_0)^e$ the field $K_{0s}(\sigma)$ is Galois over no proper subextension of $K_{0s}(\sigma)/K_0$.*

We shall also prove that if K_0 is a finitely generated infinite field and $e \geq 1$ an integer, then for almost all $\sigma \in \text{Gal}(K_0)^e$ the field $K_{0s}(\sigma)$ is a Galois extension of no proper subfield K (i.e. we remove the restriction $K_0 \subseteq K$).

It is interesting to note that if M is a minimal Henselian field, then M is a Galois extension of no proper subfield. This result follows from a theorem of Schindt, that was generalized by Engler, and later was reproved by Jarden, see [8] .

1.3. Finite PAC Extensions and the ‘Bottom Conjecture’. We classify all finite PAC extensions:

Theorem 3. *Let K/K_0 be a finite extension. Then K/K_0 is PAC if and only if one of the following holds.*

- (a) K_0 is a PAC field and K/K_0 is purely inseparable.
- (b) K_0 is real closed and K is its algebraic closure.

Let K_0 be a countable Hilbertian field. Similarly to Theorem 1, Theorem 3 can be applied to the field $K_{0s}(\sigma)$, for almost all $\sigma \in \text{Gal}(K_0)^e$. This reproves the ‘bottom theorem’ in the countable case:

Theorem 4 (Haran [11], Theorem 18.7.7). *Let K_0 be a Hilbertian field and $e \geq 1$ an integer. Then for almost all $\sigma \in \text{Gal}(K)^e$, $K_{0s}(\sigma)$ is a finite extension of no proper subfield that contains K_0 .*

Moreover, as in the case of Galois extensions, we strengthen this result and prove the ‘bottom conjecture’ [11], Problem 18.7.8 for finitely generated fields. For the precise formulation see Conjecture 7.8.

It is interesting to note that the theorems of Schmidt, Engler, and Jarden discussed before, also imply that a minimal Henselian field is a finite extension of no proper subfield.

1.4. Double Embedding Problems. Before continuing with results, we wish to briefly discuss the methods.

Arithmetic and geometric properties of a field K relate to properties of the absolute Galois group $\text{Gal}(K)$ of K . This group is equipped with the Krull topology, which makes it a profinite group. A fundamental tool in the study of profinite groups is the notion of finite embedding problems.

Here is a nice example of the above relation. A profinite group is the absolute Galois group of some PAC field if and only if all finite embedding problems for this group are weakly solvable. See Ax’ theorem [11], Theorem 11.6.2, and Lubotzky-v.d. Dries’ theorem [11], Corollary 23.1.2.

First we generalize the notion of an embedding problem for a field K to a *double embedding problem* for an extension K/K_0 . If the former is defined w.r.t. $\text{Gal}(K)$,

then the later is defined w.r.t. the restriction map $\text{Gal}(K) \rightarrow \text{Gal}(K_0)$. Roughly speaking, a double embedding problem for K/K_0 consists on two embedding problems, the ‘lower’ for K and the ‘upper’ for K_0 , which are compatible w.r.t. the restriction map.

We characterize PAC extensions in terms of special solutions of finite double embedding problems – *geometric solutions*. Those are weak solutions that are induced by some rational point of a variety, or in a different terminology, by a place of a finitely generated regular extension (see Section 2.3).

The key property that PAC extensions satisfy is the *lifting property* (Proposition 4.6). This lifting property asserts that any weak solution of the lower embedding problem can be extended to a geometric solution of the double embedding problem, provided some rationality assumption holds.

1.5. Transitivity. We prove that the PAC property of algebraic extensions is transitive.

Theorem 5. *Let $K_0 \subseteq K_1 \subseteq K_2$ be a tower of separable algebraic extensions. If both K_2/K_1 and K_1/K_0 are PAC extensions, then so is K_2/K_0 .*

This fundamental property easily follows from the lifting property. To the best of our knowledge it did not appear in the literature before. Moreover, this result together with an analog of the Ax-Lubotzky-v.d. Dries result described above leads to a new construction of PAC extensions. This will be dealt with in [1].

1.6. Descent of Galois Groups. In [22] Razon proves for a PAC extension K/K_0 that every separable extension L/K descends to a separable extension L_0/K_0 :

Theorem 6 (Razon). *Let K/K_0 be a PAC extension and L/K a separable algebraic extension. Then there exists a separable algebraic extension L_0/K_0 that is linearly disjoint from K over K_0 such that $L = L_0K$.*

We generalize Razon’s result and get the following stronger descent result.

Theorem 7. *Let K/K_0 be a PAC extension and L/K a finite Galois extension. Assume $\text{Gal}(L/K) \leq G_0$, where G_0 is regular over K_0 . Then there exists a Galois extension L_0/K_0 such that $\text{Gal}(L_0/K_0) \leq G_0$ and $L = L_0K$.*

(Here G_0 is regular over K_0 if there exists a Galois extension $F_0/K_0(t)$ with Galois group isomorphic to G_0 and such that F_0 is regular over K_0 .)

Razon’s theorem follows from Theorem 7 applied to the group $G_0 = S_n$ (for full details see the proof in Section 6). Note that the original approach of Razon to

Theorem 6 is similar to our proof but very specific: One only considers the regular realization of $G_0 = S_n$ generated by the generic polynomial $f(T_1, \dots, T_n, X) = X^n + T_1 X^{n-1} + \dots + T_n$.

Let us explain the name ‘descent’ attached to Theorem 7. If a finite Galois group $G = \text{Gal}(L/K)$ over K is regular over K_0 , then, by taking $H = G$ in Theorem 7, we get that G occurs over K_0 (since $G = \text{Gal}(L_0/K_0)$ in that case). Thus G *descends* to a Galois group over K_0 .

As a consequence of this and of the fact that finite abelian groups are regular over any field, we get, for example, that

$$K^{\text{ab}} = KK_0^{\text{ab}}.$$

Here the superscript ‘ab’ denotes the maximal abelian extension.

1.7. Acknowledgments. The author is indebted to D. Haran and M. Jarden for many discussions and suggestions concerning this work. The author thanks A. Fehm and E. Paran for valuable remarks on an earlier version of this paper, C. Meiri for his suggestion that simplified the diagram of a double embedding problem and finally the anonymous referee for his helpful comments.

2. PRELIMINARIES

In this section we define the notion of geometric solutions of an embedding problem and set up the notation and the necessary background needed for this work.

2.1. Embedding Problems. Let K be a field. Then K_s denotes its separable closure and $\tilde{K}$ its algebraic closure. The absolute Galois group of K is denoted by $\text{Gal}(K)$, i.e. $\text{Gal}(K) = \text{Gal}(K_s/K) = \text{Aut}(\tilde{K}/K)$. Recall that an **embedding problem** for $\text{Gal}(K)$ (or equivalently for K) is a diagram

$$(1) \quad \begin{array}{ccc} & & \text{Gal}(K) \\ & \nearrow \exists \theta? & \downarrow \mu \\ G & \xrightarrow{\alpha} & A, \end{array}$$

where G and A are profinite groups and μ and α are (continuous) epimorphisms. In short we write (μ, α) for (1).

A **solution** of (μ, α) is an epimorphism $\theta: \text{Gal}(K) \rightarrow G$ such that $\alpha\theta = \mu$. If θ is a homomorphism that satisfies $\alpha\theta = \mu$ but is not necessarily surjective, we say that θ is a **weak solution**. In particular, a profinite group G is a quotient of $\text{Gal}(K)$ if and only if the embedding problem $(\text{Gal}(K) \rightarrow 1, G \rightarrow 1)$ is solvable.

If G is finite (resp. α group theoretically splits), we say that the embedding problem is **finite** (resp. **split**).

The following lemma gives an obvious, but useful, criterion for a weak solution to be a solution (i.e. surjective).

Lemma 2.1. *A weak solution $\theta: \text{Gal}(K) \rightarrow G$ of an embedding problem (1) is a solution if and only if $\ker(\alpha) \leq \theta(\text{Gal}(K))$.*

Proof. Suppose $\ker(\alpha) \leq \theta(\text{Gal}(K))$. Let $g \in G$, put $a = \alpha(g)$, and let $f \in \mu^{-1}(a)$. Then $\theta(f)^{-1}g \in \ker(\alpha) \leq \theta(\text{Gal}(K))$, and hence $g \in \theta(\text{Gal}(K))$. The converse is immediate. $\square$

Two embedding problems $(\mu: \text{Gal}(K) \rightarrow A, \alpha: G \rightarrow A)$ and $(\nu: \text{Gal}(K) \rightarrow B, \beta: H \rightarrow B)$ are said to be **equivalent** if there exist isomorphisms $i: G \rightarrow H$ and $j: A \rightarrow B$ for which the following diagram commutes.

$$\begin{array}{ccccc} G & \xrightarrow{\alpha} & A & \xleftarrow{\mu} & \text{Gal}(K) \\ \downarrow i & & \downarrow j & & \parallel \\ H & \xrightarrow{\beta} & B & \xleftarrow{\nu} & \text{Gal}(K) \end{array}$$

It is evident that any (weak) solution of (μ, α) corresponds to a (weak) solution of (ν, β) and vice versa.

Denote by L the fixed field of $\ker(\mu)$ in K_s . Then μ factors as $\mu = \bar{\mu}\mu_0$, where $\mu_0: \text{Gal}(K) \rightarrow \text{Gal}(L/K)$ is the restriction map and $\bar{\mu}: \text{Gal}(L/K) \rightarrow A$ is an isomorphism. Then the embedding problems (μ, α) and $(\mu_0, \bar{\mu}^{-1}\alpha)$ are equivalent. So, from now on, we shall assume that $A = \text{Gal}(L/K)$ and μ is the restriction map (unless we explicitly specify differently).

(2)

$$\begin{array}{ccc} & \text{Gal}(K) & \\ \exists \theta? \swarrow \dots & \downarrow \mu & \\ G & \xrightarrow{\alpha} & \text{Gal}(L/K) \end{array}$$

Let $\theta: \text{Gal}(K) \rightarrow G$ be a weak solution of (μ, α) . The fixed field F of $\ker(\theta)$ is called the **solution field**. Then, if θ is a solution, the embedding problems (μ, α) and the embedding problem defined by the restriction map, i.e. $(\mu, \text{res}: \text{Gal}(F/K) \rightarrow \text{Gal}(L/K))$, are equivalent.

2.2. Geometric and Rational Embedding Problems. We define two kinds of embedding problems for a field K coming from geometric objects.

Definition 2.2. Let E be a finitely generated regular extension of K , let F/E be a Galois extension, and let $L = F \cap K_s$, where K_s is a separable closure of K . Then

the restriction map $\alpha: \text{Gal}(F/E) \rightarrow \text{Gal}(L/K)$ is surjective, since $E \cap K_s = K$. Therefore

$$(3) \quad (\mu: \text{Gal}(K) \rightarrow \text{Gal}(L/K), \alpha: \text{Gal}(F/E) \rightarrow \text{Gal}(L/K))$$

is an embedding problem for K . We call such an embedding problem **geometric embedding problem**.

If $E = K(\mathbf{t}) = K(t_1, \dots, t_e)$ is a field of rational functions over K , then we call

$$(4) \quad (\mu: \text{Gal}(K) \rightarrow \text{Gal}(L/K), \alpha: \text{Gal}(F/K(\mathbf{t})) \rightarrow \text{Gal}(L/K))$$

rational embedding problem.

We can consider only geometric embedding problems because of the following lemma.

Lemma 2.3. *Every finite embedding problem is equivalent to a geometric embedding problem.*

Proof. It follows from [11], Lemma 11.6.1. □

The following lemma shows that one can replace the tuple $\mathbf{t}$ with a single transcendental element.

Lemma 2.4. *Let K be an infinite field, $(t, \mathbf{t})$ an $(e+1)$ -tuple of variables, and (4) a rational embedding problem. Then there exists a rational embedding problem*

$$(\mu_t: \text{Gal}(K) \rightarrow \text{Gal}(L/K), \alpha_t: \text{Gal}(F_t/K(t)) \rightarrow \text{Gal}(L/K))$$

which is equivalent to (4). Furthermore, there exists a place φ of F whose residue field is F_t and such that $\varphi(t_i) = a_i + b_i t$, $a_i, b_i \in K$, and $b_i \neq 0$.

Proof. Let $a_i, b_i \in K$, $b_i \neq 0$ be as given in [4], Lemma 4, for $F/L(\mathbf{t})$. Extend the specialization $\mathbf{t} \mapsto (a_1 + b_1 t, \dots, a_e + b_e t)$ to a place of F trivial on L and let F_t be its residue field. By [4], Lemma 4, it follows that F_t is regular over L and that $[F_t : L(t)] = [F : L(\mathbf{t})]$. Then $[F_t : K(t)] = [F : K(\mathbf{t})]$, and thus $\text{Gal}(F/K(\mathbf{t})) \cong \text{Gal}(F_t/K(t))$. □

2.3. Geometric Solutions. Consider a geometric embedding problem

$$(\mu: \text{Gal}(K) \rightarrow \text{Gal}(L/K), \alpha: \text{Gal}(F/E) \rightarrow \text{Gal}(L/K))$$

for a field K . Let φ be a place of E that is unramified in F . We always assume that φ is trivial on K , i.e. $\varphi(x) = x$, for $x \in K$. We denote a residue field by bar, e.g., $\bar{E}$ is the residue field of E .

Assume φ is K -rational, i.e. $\bar{E} = K$, and extend it to a place of F , say Φ . By composing Φ with an appropriate Galois automorphism, we can assume that Φ is trivial on L .

Then Φ/φ canonically induces a weak solution Φ^* of (μ, α) . The image of Φ^* is the decomposition group and $\Phi(\Phi^*(\sigma)(x)) = \sigma(\Phi(x))$ for all $\sigma \in \text{Gal}(K)$, provided $\Phi(x)$ is finite. If we choose a different extension of φ to F , say Ψ , then Ψ^* and Φ^* differ by an inner automorphism of $\text{Gal}(F/E)$. For complete details see [11], Lemma 6.1.4).

Definition 2.5. Let $\theta: \text{Gal}(K) \rightarrow \text{Gal}(F/E)$ be a weak solution of (μ, α) . Then we call it **geometric** if there exists a place Φ of F unramified over E such that $\bar{E} = K$ and $\theta = \Phi^*$.

If we have a commutative diagram

$$\begin{array}{ccc} & & \text{Gal}(K) \\ & & \downarrow \mu_1 \\ H_1 & \xrightarrow{\alpha_1} & G_1 \\ \downarrow \pi & & \downarrow \mu_2 \\ H_2 & \xrightarrow{\alpha_2} & G_2 \end{array}$$

then we say that (μ_1, α_1) **dominates** (μ_2, α_2) . Note that any (weak) solution $\theta_1: \text{Gal}(K) \rightarrow H_1$ of (μ_1, α_1) induces a (weak) solution $\theta_2 = \pi\theta_1$ of (μ_2, α_2) .

Geometric solutions are compatible with scalar extensions:

Lemma 2.6. *Consider a geometric embedding problem*

$$(\mu: \text{Gal}(K) \rightarrow \text{Gal}(L/K), \alpha: \text{Gal}(F/E) \rightarrow \text{Gal}(L/K)).$$

Let M/K be a Galois extension with $L \subseteq M$. Then the geometric embedding problem

$$(\mu': \text{Gal}(K) \rightarrow \text{Gal}(M/K), \alpha': \text{Gal}(FM/E) \rightarrow \text{Gal}(M/K)),$$

where α' and μ' are the corresponding restriction maps, dominates the embedding problem (μ, α) with respect to the restriction maps. Furthermore, if Ψ^ is a geometric (weak) solution of (μ', α') , then $(\Psi|_F)^*$ is a geometric (weak) solution of (μ, α) .*

Proof. As E/K is regular, we have

$$\begin{aligned} \text{Gal}(FN/E) &= \text{Gal}(F/E) \times_{\text{Gal}(LE/E)} \text{Gal}(ME/E) \\ &\cong \text{Gal}(F/E) \times_{\text{Gal}(L/K)} \text{Gal}(M/K) \end{aligned}$$

and the projection maps coincide with the restriction maps. Thus (μ', α') dominates (μ, α) . Now let Ψ^* be a geometric (weak) solution of (μ', α') . For $\Phi = \Psi|_F$, we have that Φ is unramified over E and $\text{res}_{FM,F} \circ \Psi^* = \Phi^*$, as needed. $\square$

3. PSEUDO ALGEBRAICALLY CLOSED EXTENSIONS AND DOUBLE EMBEDDING PROBLEMS

3.1. Basic Properties. The following proposition gives several equivalent definitions of PAC extensions in terms of polynomials and places, including a reduction to plane curves. A proof of that proposition essentially appears in [15]. Nevertheless, for the sake of completeness, we give here a formal proof.

Proposition 3.1. *The following conditions are equivalent for a field extension K/K_0 .*

- (3.1a) K/K_0 is PAC.
- (3.1b) For every $e \geq 1$ and every absolutely irreducible polynomial $f(\mathbf{T}, X) \in K[T_1, \dots, T_e, X]$ that is separable in X , and nonzero $r(\mathbf{T}) \in K[\mathbf{T}]$ there exists $(\mathbf{a}, b) \in K_0^e \times K$ for which $r(\mathbf{a}) \neq 0$ and $f(\mathbf{a}, b) = 0$.
- (3.1c) For every absolutely irreducible polynomial $f(T, X) \in K[T, X]$ that is separable in X , and nonzero $r(T) \in K(T)$ there exists $(a, b) \in K_0 \times K$ for which $r(a) \neq 0$ and $f(a, b) = 0$.
- (3.1d) For every finitely generated regular extension E/K with separating transcendence basis $\mathbf{t} = (t_1, \dots, t_e)$ and every nonzero $r(\mathbf{t}) \in K(\mathbf{t})$, there exists a K -rational place φ of E unramified over $K(\mathbf{t})$ such that $\overline{K_0(\mathbf{t})} = K_0$, $\mathbf{a} = \varphi(\mathbf{t})$ is finite, and $r(\mathbf{a}) \neq 0, \infty$.
- (3.1e) For every finitely generated regular extension E/K with separating transcendence basis t and every nonzero $r(t) \in K[t]$ there exists a K -rational place φ of E unramified over $K(t)$ such that $\overline{K_0(t)} = K_0$, $a = \varphi(t) \neq \infty$, and $r(a) \neq 0, \infty$.

Proof. The proof of [15], Lemma 1.3, gives the equivalence between (3.1a), (3.1b), and (3.1c). Obviously (3.1d) implies (3.1e), so it suffices to prove that (3.1b) implies (3.1d) and that (3.1e) implies (3.1c).

(3.1b) $\Rightarrow$ (3.1d): Let $x \in E/K(\mathbf{t})$ be integral over $K[\mathbf{t}]$ such that $E = K(\mathbf{t}, x)$. Let $f(\mathbf{T}, X) \in K[\mathbf{T}, X]$ be the absolutely irreducible polynomial which is monic and separable in X and for which $f(\mathbf{t}, x) = 0$. Let $0 \neq g(\mathbf{t}) \in K[\mathbf{t}]$ be the discriminant of $f(\mathbf{T}, X)$ as a polynomial in X . We have $(\mathbf{a}, b) \in K_0^e \times K$ such that $f(\mathbf{a}, b) = 0$ and $g(\mathbf{a})r(\mathbf{a}) \neq 0, \infty$. Extend the specialization $\mathbf{t} \mapsto \mathbf{a}$ to a K -rational place φ of E

with the following properties to conclude the implication: (1) $\varphi(x) = b \neq \infty$ (this is possible since x is integral over $K[t]$); (2) $\overline{K_0(\mathbf{t})} = K_0$ and $\bar{E} = K(b) = K$ ([11], Lemma 2.2.7.); (3) φ is unramified over $K(\mathbf{t})$ ([11], Lemma 6.1.8).

(3.1e) $\Rightarrow$ (3.1c): Let $f(T, X) = \sum_{k=0}^n a_k(T)X^k$ and $r(T)$ be as in (3.1c). Set $r'(T) = r(T)a_n(T)$. Let t be a transcendental element and let $x \in \widehat{K(t)}$ be such that $f(t, x) = 0$. Let $E = K(t, x)$. Then E is regular over K and separable over $K(t)$. Applying (3.1e) to E and $r'(t)$ we get a K -rational place φ of E satisfying the following properties. (1) $a = \varphi(t) \in K_0$ which implies that $b = \varphi(x)$ is finite, since $\varphi(f(t, x)) = 0$ and $f(a, X)$ has a nonzero leading coefficient; (2) $\bar{E} = K$, which concludes the proof since $b \in \bar{E} = K$. $\square$

3.2. Geometric Solutions and PAC fields. The following result characterizes when a solution is geometric in terms of a rational place of some regular extension. This sharpens earlier works of Roquette on PAC Hilbertian fields [11], Corollary 27.3.3, and of Fried-Haran-Jarden on Frobenius fields [11], Proposition 24.1.4. In the case where $L = K$, it was also proved by Dèbes in his work on the Beckmann-Black problem [7].

Proposition 3.2. *Let K be a field and consider a geometric embedding problem*


$$(\mu: \text{Gal}(K) \rightarrow \text{Gal}(L/K), \alpha: \text{Gal}(F/E) \rightarrow \text{Gal}(L/K))$$

for K . Let $\theta: \text{Gal}(K) \rightarrow \text{Gal}(F/E)$ be a weak solution. Then there exists a finite separable extension $\hat{E}/E$ such that $\hat{E}/K$ is regular and for every place φ of E/K that is unramified in F the following two conditions are equivalent.

- (1) φ extends to a place Φ of F such that $\bar{E} = K$ and $\Phi^* = \theta$.
- (2) φ extends to a K -rational place of $\hat{E}$.

Proof. First we consider the special case when $\text{Gal}(F/E) \cong \text{Gal}(L/K)$. Then there is a unique solution of (μ, α) , namely $\theta = \alpha^{-1}\mu$. Let Φ be an extension of φ to F and take $\hat{E} = E$. It is trivial that (1) implies (2). Assume (2). Then Φ^* is defined, and from the uniqueness, $\Phi^* = \theta$.

Next we prove the general case. Let M/K be a Galois extension such that $\text{Gal}(M) = \ker(\theta)$ (in particular, $L \subseteq M$) and let $\hat{F} = FM$.



As F and M are linearly disjoint over L , the fields F and EM are linearly disjoint over EL . We have

$$\mathrm{Gal}(\hat{F}/E) = \mathrm{Gal}(F/E) \times_{\mathrm{Gal}(L/K)} \mathrm{Gal}(M/K).$$

Define $\hat{\theta}: \mathrm{Gal}(K) \rightarrow \mathrm{Gal}(\hat{F}/E)$ by $\hat{\theta}(\sigma) = (\theta(\sigma), \sigma|_M)$. Let $\hat{E}$ denote the fixed field of $\hat{\theta}(\mathrm{Gal}(K))$ in $\hat{F}$. Then $\hat{\theta}$ is a solution in

$$\begin{array}{ccc} & \mathrm{Gal}(K) & \\ \hat{\theta} \swarrow & \downarrow \mu & \\ \mathrm{Gal}(\hat{F}/\hat{E}) & \xrightarrow{\hat{\alpha}} & \mathrm{Gal}(L/K). \end{array}$$

Here $\hat{\alpha}$ is the restriction map. In particular, $\hat{E}/K$ is regular. Also $\ker(\hat{\theta}) = \ker(\theta) \cap \mathrm{Gal}(M) = \mathrm{Gal}(M)$, so $\mathrm{Gal}(\hat{F}/\hat{E}) \cong \mathrm{Gal}(M/K)$.

Assume φ extend to a K -rational place $\hat{\varphi}$ of $\hat{E}$. Extend $\hat{\varphi}$ L -linearly to a place $\hat{\Phi}$ of $\hat{F}$. Then $\hat{\Phi}/\hat{\varphi}$ is unramified. Let $\Phi = \hat{\Phi}|_F$. Then by the first part $\hat{\Phi}^* = \hat{\theta}$. Lemma 2.6 then asserts that $\Phi^* = \theta$.

On the other hand, assume that φ extends to a place Φ of F such that $\bar{E} = K$ and $\Phi^* = \theta$. Extend Φ M -linearly to a place $\hat{\Phi}$ of $\hat{F}$. Then, since $\mathrm{res}_{\hat{F},F}(\hat{\Phi}^*) = \Phi^*$ and $\mathrm{res}_{\hat{F},M}(\hat{\Phi}^*) = \mathrm{res}_{K_s,M}$, we have

$$\hat{\Phi}^*(\sigma) = (\Phi^*(\sigma), \sigma|_M) = \hat{\theta}.$$

Therefore the residue field of $\hat{E}$ is also K . □

Remark 3.3. In the proof it was shown that $\hat{E} \subseteq FM$, where M is the solution field of θ .

Proposition 3.2 is extremely useful. We first apply it to PAC fields.

Corollary 3.4. *Every weak solution of a finite geometric embedding problem for a PAC field K is geometric.*

Proof. Let $(\mu: \mathrm{Gal}(K) \rightarrow \mathrm{Gal}(L/K), \alpha: \mathrm{Gal}(F/E) \rightarrow \mathrm{Gal}(L/K))$ be a finite geometric embedding problem for K and θ a solution. Let $\hat{E}$ be the regular extension of K given in Proposition 3.2. There exists a K -rational place $\hat{\varphi}$ of $\hat{E}$. Then by the proposition we can extend $\hat{\varphi}|_E$ to a place Φ of F such that $\Phi^* = \theta$. □

3.3. Double Embedding Problems. In this section we generalize the notion of embedding problems to field extensions.

3.3.1. *The Definition of Double Embedding Problems.* Let K/K_0 be a field extension. A **double embedding problem (DEP)** for K/K_0 consists of two embedding problems: $(\mu: \text{Gal}(K) \rightarrow G, \alpha: H \rightarrow G)$ for K and $(\mu_0: \text{Gal}(K_0) \rightarrow G_0, \alpha_0: H_0 \rightarrow G_0)$ for K_0 , which are compatible in the following sense. $H \leq H_0$, $G \leq G_0$, and if we write $i: H \rightarrow H_0$ and $j: G \rightarrow G_0$ for the inclusion maps and r for the restriction map $\text{Gal}(K) \rightarrow \text{Gal}(K_0)$, then the following diagram commutes.

$$(5) \quad \begin{array}{ccccc} & & \text{Gal}(K) & & \\ & \swarrow \exists \theta? & \downarrow r & \searrow & \\ & & \text{Gal}(K_0)_\mu & & \\ & \swarrow \exists \theta_0? & \downarrow \mu_0 & \searrow & \\ H & \xrightarrow{i} & H_0 & \xrightarrow{\alpha_0} & G_0 & \xleftarrow{j} & G \\ & \searrow \alpha & & & & & \end{array}$$

Given a DEP for K/K_0 , we refer to the corresponding embedding problem for K (resp. K_0) as the **lower** (resp. the **upper**) embedding problem. We call a DEP **finite** if the upper (and hence also the lower) embedding problem is finite.

A **weak solution** of a DEP (5) is a weak solution θ_0 of the upper embedding problem which restricts to a weak solution θ of the lower embedding problem via the restriction map $r: \text{Gal}(K) \rightarrow \text{Gal}(K_0)$. In case K/K_0 is a separable algebraic extension, the restriction map is the inclusion map, and hence the condition on θ_0 reduces to $\theta_0(\text{Gal}(K)) \leq H$. To emphasize the existence of θ , we usually regard a weak solution of a DEP as a pair (θ, θ_0) (where θ is the restriction of θ_0 to $\text{Gal}(K)$).

3.3.2. *Rational Double Embedding Problems.* Consider a double embedding problem (5) and let L_0 and L be the fixed fields of the kernels of μ_0 and μ , respectively. Then we have isomorphisms $\bar{\mu}_0: \text{Gal}(L_0/K_0) \rightarrow G_0$ and $\bar{\mu}: \text{Gal}(L/K) \rightarrow G$. Hence (as in the case of embedding problems) replacing G_0 and G with $\text{Gal}(L_0/K_0)$ and $\text{Gal}(L/K)$ (and replacing correspondingly all the maps) gives us an equivalent DEP. The compatibility condition is realized as $L = L_0K$.

In the context of this work we are mainly interested in double embedding problems which satisfy some rationality condition.

Definition 3.5. If in a double embedding problem the upper embedding problem is rational, we say that the double embedding problem is **rational**.

Lemma 3.6. *If (5) is a rational DEP, then the upper embedding problem is rational and the lower embedding problem is geometric.*

Moreover, we can take $e = 1$, i.e. $\mathbf{t} = t$ – a transcendental element.

Proof. Let (5) be a rational double embedding problem. By definition, it means that $G_0 = \text{Gal}(L_0/K_0)$, there exists a regular extension F_0 of L_0 and a separating transcendence basis $\mathbf{t} \in F_0$ such that $F_0/K_0(\mathbf{t})$ is Galois with Galois group $H_0 = \text{Gal}(F_0/K_0(\mathbf{t}))$, and $\alpha_0: H_0 \rightarrow G_0$ is the restriction map. By Lemma 2.4 we may assume that $\mathbf{t} = t$.

Let $F = F_0K$ and $L = L_0K$. The compatibility condition implies that H embeds into $\text{Gal}(F_0/K_0(\mathbf{t}))$ (via i) as a subgroup of $\text{Gal}(F_0/L_0 \cap K(\mathbf{t})) \cong \text{Gal}(F/K(\mathbf{t}))$. Let $E \subseteq F$ be the fixed field of H , i.e., $H = \text{Gal}(F/E)$. Under this embedding, $\alpha: \text{Gal}(F/K(\mathbf{t})) \rightarrow \text{Gal}(L/K)$ is the restriction map. Therefore $\alpha(H) = \text{Gal}(L/K)$ implies that $E \cap L = K$, and hence E is regular over K .

$$(6) \quad \begin{array}{ccccc} & & \text{Gal}(K) & & \\ & & \downarrow r & \searrow \mu & \\ & & \text{Gal}(K_0) & & \\ & & \downarrow \mu_0 & & \\ \text{Gal}(F/E) & \xhookrightarrow{i} & \text{Gal}(F_0/K_0(t)) & \xrightarrow{\alpha_0} & \text{Gal}(L_0/K_0) & \xleftarrow{j} & \text{Gal}(L/K). \\ & & \searrow \alpha & & \end{array}$$

Consequently the lower embedding problem is geometric. $\square$

Remark 3.7. The converse of the above lemma is also valid, that is to say, assume we have a rational double embedding problem as in (6), i.e. a finitely generated regular extension E/K , a separating transcendence basis $\mathbf{t}$ for E/K , and a finite Galois extension $F_0/K_0(\mathbf{t})$ such that $E \subseteq F$, where $F = F_0K$. Then all the restriction maps in (6) are surjective, and hence (6) defines a finite double embedding problem.

3.3.3. Geometric Solutions of Double Embedding Problems. First recall that a weak solution θ of an embedding problem

$$(\mu: \text{Gal}(K) \rightarrow \text{Gal}(L/K), \alpha: \text{Gal}(F/E) \rightarrow \text{Gal}(L/K))$$

is geometric if $\theta = \Phi^*$, for some place Φ of F that is unramified over E and under which the residue field of E is K . Then we call a weak solution (θ, θ_0) of (6) **geometric** if $(\theta, \theta_0) = (\Phi^*, \Phi_0^*)$, where Φ^* is a geometric solution of the lower embedding problem and $\Phi_0 = \Phi|_{F_0}$.

Note that since Φ_0^* is a solution of the upper embedding problem, the residue field of $K_0(\mathbf{t})$ is K_0 . In particular, if $\Phi(\mathbf{t})$ is finite, then $\Phi(\mathbf{t}) \in K_0^e$. Also note that

for a place of F that is unramified over E and such that $\bar{E} = K$ and $\overline{K_0(\mathbf{t})} = K_0$, the pair (Φ^*, Φ_0^*) is indeed a weak solution of (6), since $\Phi_0^* = \text{res}_{K_s, K_{0s}} \Phi^*$.

4. THE LIFTING PROPERTY

In this section we formulate and prove the lifting property. First we reduce the discussion to separable algebraic extensions by showing that if K/K_0 is PAC, then $K \cap K_{0s}/K_0$ is PAC and $\text{Gal}(K) \cong \text{Gal}(K \cap K_{0s})$ via the restriction map. Then we characterize separable algebraic PAC extensions in terms of geometric solutions of double embedding problems. From this characterization we establish the lifting property. Finally we prove a strong (but complicated) version of the lifting problem to PAC extensions of finitely generated fields.

4.1. Reduction to Separable Algebraic Extensions. In [15], Corollary 1.5, Jarden and Razan show

Lemma 4.1 (Jarden-Razon). *If K/K_0 is PAC, then so is $K \cap K_{0s}/K_0$.*

Moreover, we have

Theorem 4.2. *Let K/K_0 be a PAC extension. Then $K \cap K_{0s}/K_0$ is PAC and the restriction map $\text{Gal}(K) \rightarrow \text{Gal}(K \cap K_{0s})$ is an isomorphism.*

Proof. It suffices to show that $K_s = K_{0s}K$. Let L/K be a finite Galois extension with Galois group G of order n . Embed G into the symmetric group S_n . Let $F_0/K_0(\mathbf{t})$ be a regular realization of S_n with F_0 algebraically independent from K over K_0 ([18], p. 272, Example 4). Then $F = F_0K$ is regular over K and $\text{Gal}(F/K(\mathbf{t})) \cong S_n$. Furthermore

$$\begin{aligned} \text{Gal}(FL/K(\mathbf{t})) &= \text{Gal}(FL/L(\mathbf{t})) \times \text{Gal}(FL/F) \\ &\cong \text{Gal}(F/K(\mathbf{t})) \times \text{Gal}(L/K) \cong S_n \times G. \end{aligned}$$

Let E be the fixed field of the subgroup $\Delta = \{(g, g) \mid g \in G\}$ in FL , i.e., $\text{Gal}(FL/E) \cong \Delta$. By Galois correspondence, $S_n \Delta = S_n \times G$ implies that $E \cap L = K$ and $1 = \Delta \cap S_n = G \cap \Delta$ implies that $FL = EL = FE$. In particular, E/K is regular.

$$\begin{array}{ccc} F & \text{---} & FL \\ | & \nearrow E & | \\ K(\mathbf{t}) & \text{---} & L(\mathbf{t}) \\ | & & | \\ K & \text{---} & L \end{array}$$

As K/K_0 is PAC, there is a K -rational place φ of E such that $\overline{K_0(\mathbf{t})} = K_0$, $\mathbf{a} = \varphi(\mathbf{t})$ is finite, and $h(\mathbf{a}) \neq 0$. Extend φ to a place Φ of FL which is trivial on L .

Then since Φ/φ is unramified, $FL = EL$ implies that $\overline{FL} = \overline{EL} = L$ ([11], Lemma 2.4.8). However, as $F = F_0K$, it follows that $\bar{F} = \bar{F}_0K$, hence (again by [11], Lemma 2.4.8), $L = \overline{FL} = \overline{FE} = \bar{F} = \bar{F}_0K \subseteq K_{0s}K$, as needed. $\square$

Remark 4.3. The above theorem also follows from Razon's result Theorem 6. However we shall prove the converse implication.

PAC fields have a nice elementary theory. Since K and $K \cap K_{0s}$ are PAC fields and since they have isomorphic absolute Galois groups, they are elementary equivalent under some necessary condition:

Corollary 4.4. *Let K/K_0 be a separable PAC extension. Assume that K and $K \cap K_{0s}$ have the same degree of imperfection. Then K is an elementary extension of $K \cap K_{0s}$.*

Proof. The assertion follows from [11], Corollary 20.3.4, and Theorem 4.2. $\square$

4.2. Characterization of Separable Algebraic PAC Extensions.

Proposition 4.5. *Let K/K_0 be a separable algebraic field extension. The following conditions are equivalent:*


- (4.5a) K/K_0 is PAC.
- (4.5b) *For every finite rational double embedding problem (6) for K/K_0 and every nonzero rational function $r(\mathbf{t}) \in K(\mathbf{t})$, there exists a geometric weak solution (Φ^*, Φ_0^*) such that $\mathbf{a} = \Phi(\mathbf{t})$ is finite and $r(\mathbf{a}) \neq 0, \infty$.*
- (4.5c) *For every finite rational double embedding problem (6) for K/K_0 with $\mathbf{t} = t$ a transcendental element there exist infinitely many geometric weak solutions (Φ^*, Φ_0^*) .*

Proof. The implication (4.5a) $\Rightarrow$ (4.5b) follows from Proposition 3.1 (part (3.1d)) and the definition of geometric weak solutions.

Taking $\mathbf{t} = t$ a transcendental element (instead of a general tuple) in (4.5c) yields (4.5b).

(4.5c) $\Rightarrow$ (4.5a): We apply Proposition 3.1 and show that (3.1e) holds. Let E/K be a regular extension with a separating transcendence basis t and let $r(t) \in K[t]$ be nonzero. Choose F_0 to be a finite Galois extension of $K_0(t)$ such that $E \subseteq F_0K$ (such F_0 exists since K/K_0 is separable and algebraic). Let $F = F_0K$, $L = F \cap K_s$, and

$L_0 = F_0 \cap K_{0s}$. By assumption there are infinitely many geometric weak solutions (Φ^*, Φ_0^*) of the DEP



Since for only finitely many solutions $\Phi(t)$ is infinite or $r(\Phi(t)) = 0, \infty$ we can find a solution such that $\Phi(t) \neq \infty$ and $r(\Phi(t)) \neq 0, \infty$. In particular, $\bar{E} = K$ and $\overline{K_0(t)} = K_0$, as required in (3.1e). $\square$

Let K/K_0 be a PAC extension and consider a rational DEP for K/K_0 . The following key property – the lifting property – asserts that any weak solution of the lower embedding problem can be lifted to a geometric weak solution of the DEP.

Proposition 4.6 (The lifting property). *Let K/K_0 be a PAC extension, let (6) be a rational DEP for K/K_0 , and let $\theta: \text{Gal}(K) \rightarrow \text{Gal}(F/E)$ be a weak solution of the lower embedding problem in (6). Then there exists a geometric weak solution (Φ^*, Φ_0^*) of (6) such that $\theta = \Phi^*$.*

Moreover, if $r(\mathbf{t}) \in K(\mathbf{t})$ is nonzero, we can choose Φ such that $\mathbf{a} = \Phi(\mathbf{t}) \in K_0^e$ and $r(\mathbf{a}) \neq 0, \infty$.

Proof. By Proposition 3.2 there exists a finite separable extension $\hat{E}/E$ that is regular over K with the following property. If a place φ of E that is unramified in F can be extended to a K -rational place of $\hat{E}$, then it can be extended to a place Φ of F such that $\Phi^* = \theta$.

By the PACness of K/K_0 there exists a K -rational place $\hat{\varphi}$ of $\hat{E}$ such that $\varphi = \hat{\varphi}|_E$ is unramified in F , the residue field of $K_0(\mathbf{t})$ is K_0 , $\mathbf{a} = \Phi(\mathbf{t})$ is finite, and $r(\mathbf{a}) \neq 0, \infty$. If we extend φ to the place Φ of F given above and let $\Phi_0 = \varphi|_{F_0}$, then we get that (Φ^*, Φ_0^*) is a geometric weak solution and that $\Phi^* = \theta$. $\square$

The first easy consequence of the lifting property is the transitivity of PAC extensions. We first prove Theorem 5 and then deduce a more general result.

Proof of Theorem 5. Let $K_0 \subseteq K_1 \subseteq K_2$ be a tower of separable algebraic extensions and assume that K_2/K_1 and K_1/K_0 are PAC extensions. We need to prove that K_2/K_0 is PAC too.

Let

$$((\mu_0: \text{Gal}(K_0) \rightarrow \text{Gal}(L_0/K_0), \alpha_0: \text{Gal}(F_0/K_0(t)) \rightarrow \text{Gal}(L_0/K_0)), \\ (\mu_2: \text{Gal}(K_2) \rightarrow \text{Gal}(L_2/K_2), \alpha_2: \text{Gal}(F_2/E) \rightarrow \text{Gal}(L_2/K_2)))$$

be a rational finite DEP for K_2/K_0 . By Lemma 4.5 it suffices to find a geometric weak solution to $((\mu_0, \alpha_0), (\mu_2, \alpha_2))$. Set $F_1 = F_0K_1$, $L_1 = L_0K_1$. Then, since K_2/K_1 is PAC there exists a weak solution (Φ_2^*, Φ_1^*) of the double embedding problem defined by the lower part of the following commutative diagram.

(7)

$$\begin{array}{ccccc}
 & & \text{Gal}(K_0) & & \\
 & \swarrow \Phi_0^* & \uparrow & \searrow \mu_0 & \\
 \text{Gal}(F_0/K_0(t)) & \xrightarrow{\alpha_0} & & & \text{Gal}(L_0/K_0) \\
 \uparrow & & \text{Gal}(K_1) & & \uparrow \\
 & \swarrow \Phi_1^* & \uparrow & \searrow \mu_1 & \\
 \text{Gal}(F_1/K_1(t)) & \xrightarrow{\alpha_1} & & & \text{Gal}(L_1/K_1) \\
 \uparrow & & \text{Gal}(K_2) & & \uparrow \\
 & \swarrow \Phi_2^* & \uparrow & \searrow \mu_2 & \\
 \text{Gal}(F_2/E) & \xrightarrow{\alpha_2} & & & \text{Gal}(L_2/K_2).
 \end{array}$$

Now we lift Φ_1^* to a geometric weak solution (Φ_1^*, Φ_0^*) of the DEP for K_1/K_0 defined by the higher part of the diagram. This is possible by the lifting property applied to the PAC extension K_1/K_0 .

Since $\Phi_0^*|_{\text{Gal}(K_2)} = \Phi_1^*|_{\text{Gal}(K_2)} = \Phi_2^*$ we get that (Φ_2^*, Φ_0^*) is a geometric weak solution of the DEP we started from. $\square$

Corollary 4.7. *Let κ be an ordinal number and let*

$$K_0 \subseteq K_1 \subseteq K_2 \subseteq \cdots \subseteq K_\kappa$$

be a tower of separable algebraic extensions. Assume that $K_{\alpha+1}/K_\alpha$ is PAC for every $\alpha < \kappa$ and that $K_\alpha = \bigcup_{\beta < \alpha} K_\beta$ for every limit $\alpha \leq \kappa$. Then K_κ/K_0 is PAC.

Proof. We apply transfinite induction. Let $\alpha \leq \kappa$. If α is a successor ordinal, then the assertion follows from Theorem 5.

Let α be a limit ordinal. Consider a rational DEP

$$(8) \quad \begin{array}{ccccc} & & \text{Gal}(K_\alpha) & & \\ & & \downarrow & \searrow & \\ & & \text{Gal}(K_0) & & \\ & & \downarrow & & \\ \text{Gal}(F_\alpha) & \longrightarrow & \text{Gal}(F_0/K_0(\mathbf{t})) & \longrightarrow & \text{Gal}(L_0/K_0) \longleftarrow \text{Gal}(L_\alpha/K_\alpha) \\ & \searrow & & \nearrow & \end{array}$$

Here $F_\alpha = F_0K_\alpha$ and $L_\alpha = L_0K_\alpha$. Now since all the extensions are finite, there exists $\beta < \alpha$ such that $\text{Gal}(F_\alpha/K_\alpha) \cong \text{Gal}(F_\beta/K_\beta)$ and $\text{Gal}(L_\alpha/K_\alpha) \cong \text{Gal}(L_\beta/K_\beta)$ (via the corresponding restriction maps), where $F_\beta = F_0K_\beta$ and $L_\beta = L_0K_\beta$.

Induction gives a weak solution of the double embedding problem (8) with β replacing α . This weak solution induces a weak solution of (8) via the above isomorphisms. $\square$

4.3. Strong Lifting Property for PAC Extensions of Finitely Generated Fields. Let K_0 be a finitely generated field (over its prime field). In this section we prove a strong lifting property for PAC extensions K/K_0 . The additional ingredient is the Mordell conjecture for finitely generated fields (now a theorem due to Faltings in characteristic 0 [9] and to Grauert-Manin in positive characteristic [24], page 107).

The following lemma is based on the Mordell conjecture.

Lemma 4.8 ([15], Proposition 5.4). *Let K_0 be a finitely generated infinite field, $f \in K_0[T, X]$ an absolutely irreducible polynomial which is separable in X , $g \in K_0[T, X]$ an irreducible polynomial which is separable in X , and $0 \neq r \in K_0[T]$. Then there exist a finite purely inseparable extension K'_0 of K_0 , a nonconstant rational function $q \in K'_0(T)$, and a finite subset B of K'_0 such that $f(q(T), X)$ is absolutely irreducible, $g(q(a), X)$ is irreducible in $K'_0[X]$, and $r(q(a)) \neq 0$ for any $a \in K'_0 \setminus B$.*

Let K/K_0 be an extension. Consider a rational double embedding problem (6) for K/K_0 (with $\mathbf{t} = t$). For any subextension K_1 of K/K_0 we have a corresponding rational double embedding problem. Namely

$$(9) \quad \begin{array}{ccccc} & & \text{Gal}(K) & & \\ & & \downarrow & \searrow \mu & \\ & & \text{Gal}(K_1) & & \\ & & \downarrow \mu_1 & & \\ \text{Gal}(F/E) & \xrightarrow{\quad} & \text{Gal}(F_1/K_1(t)) & \xrightarrow{\alpha_1} & \text{Gal}(L_1/K_1) & \xleftarrow{\quad} & \text{Gal}(L/K), \\ & \searrow \alpha & & & & & \end{array}$$

where $F_1 = F_0K_1$, $L_1 = L_0K_1$, and μ_1 and α_1 are the restriction maps.

Assume that K'_1/K_1 is a purely inseparable extension. Then the double embedding problem (9) remains the same if we replace all fields by their compositum with K'_1 .

Proposition 4.9 (Strong lifting property). *Let K be a PAC extension of a finitely generated field K_0 . Let*

$$\mathcal{E}(K) = (\mu: \text{Gal}(K) \rightarrow \text{Gal}(L/K), \alpha: \text{Gal}(F/E) \rightarrow \text{Gal}(L/K))$$

be a geometric embedding problem and let $\theta: \text{Gal}(K) \rightarrow \text{Gal}(F/E)$ be a weak solution of $\mathcal{E}(K)$. Then there exist a finite subextension K_1/K_0 and a finite purely inseparable extension K'_1/K_1 satisfying the following properties.

- (a) *For any rational double embedding problem (9) for K/K_1 whose lower embedding problem is $\mathcal{E}(K)$, we can lift θ to a weak solution (θ, θ_1) of the double embedding problem (9) in such a way that θ_1 is surjective.*
- (b) *The solution (θ, θ_1) is a geometric solution of the double embedding problem that we get from (9) by replacing all fields with their compositum with K'_1 .*

Proof. By Proposition 3.2 there exists a finite separable $\hat{E}/E$ that is regular over K such that a K -place φ of E that is unramified in F satisfies $\varphi^* = \theta$ if and only if φ extends to a K -rational place of $\hat{E}$. Let $f(t, X) \in K[t, X]$ be an absolutely irreducible polynomial whose root x generates $\hat{E}/K(t)$, i.e. $\hat{E} = K(t, x)$. Let M be the fixed field of $\ker(\theta)$ in K_s . Then M/K is a finite Galois extension. Let $h(X) \in K[X]$ be a Galois irreducible polynomial whose root generates M/K .

Let K_1 be a finite subextension of K/K_0 that contains the coefficients of f and h and such that h is Galois over it. Let M_1 be the splitting field of h over K_1 and let L_1, F_1 be as in the corresponding rational double embedding problem (9). Then $\text{Gal}(M/K) \cong \text{Gal}(M_1/K_1)$, and thus also $\text{Gal}(L/K) \cong \text{Gal}(L_1/K_1)$.

$$\begin{array}{ccccccc}
& & & & F & & \\
& & & & | & & \\
E & \text{---} & EL & \text{---} & EM & & \\
| & & | & & | & & \\
K(t) & \text{---} & L(t) & \text{---} & M(t) & & \\
| & & | & & | & & \\
& & & & F_1 & & \\
& & & & | & & \\
K_0(t) & \text{---} & K_1(t) & \text{---} & L_1(t) & \text{---} & M_1(t)
\end{array}$$

Let $g(t, X) \in K_1[T, X]$ be an irreducible polynomial whose root generates $F_1/K_1(t)$. Choose $r(t) \in K_1(t)$ such that $r(a) \neq 0$ implies that the prime $(t - a)$ is unramified in F_1 and that the leading coefficients of $f(t, X)$ and $g(t, X)$ do not vanish at a . Let K'_1/K_1 be the purely inseparable extension, $B \subseteq K'_1$ the finite subset, and $q \in K'_1(T)$ the nonconstant rational function that Lemma 4.8 gives for K_1 , g , f , and r . Let $K' = KK'_1$.

Since K'/K'_1 is PAC ([15], Corollary 2.5) there exist $a \in K'_1 \setminus B$ and $b \in K'$ for which $f(q(a), b) = 0$ (Proposition 3.1). Extend $t \mapsto q(a)$ to a K' -rational place $\hat{\varphi}$ of $\hat{E}K'_1$. Then $\varphi = \hat{\varphi}|_{EK'_1}$ is unramified in FK'_1 (since $r(q(a)) \neq 0$).

By Proposition 3.2 φ extends to a place Φ of FK'_1 such that $\Phi^* = \theta$ and write $\Phi_1 = \Phi|_{F_1}$. Then (Φ^*, Φ_1^*) is a geometric weak solution of the DEP $((\mu, \alpha), (\mu_1, \alpha_1))$ that we get from (9) by replacing all fields with their compositum with K'_1 .

Moreover, since $F_1 K'_1 / K'_1(t)$ is generated by $g(t, X)$ and $g(q(a), X)$ is irreducible, we get that Φ_1^* is surjective. This proves (b). Now assertion (a) follows since (Φ^*, Φ_1^*) is a (not necessarily geometric) solution of $((\mu, \alpha), (\mu_1, \alpha_1))$. $\square$

5. THE GALOIS CLOSURE OF PAC EXTENSIONS

This section proves the main result of the paper, Theorem 1, that says the Galois closure of a proper separable algebraic PAC extension is the separable closure. The proof uses the lifting property and some properties of realizing wreath products in fields. We start by recalling the latter and then prove the theorem.

5.1. Wreath Products in Fields. Let A and G be finite groups. The **wreath product** $A \wr G$ is defined to be the semidirect product $A^G \rtimes G$, where G acts on A^G by translation. More precisely, $A^G = \{f: G \rightarrow A\}$ and

$$f^\sigma(\tau) = f(\sigma\tau)$$

for all $f \in A^G$ and $\sigma, \tau \in G$. Then each element of $A \wr G$ can be written uniquely as $f\sigma$, $f \in A^G$ and $\sigma \in G$ and the multiplication is given by

$$(f\sigma)(g\tau) = fg^{\sigma^{-1}}\sigma\tau.$$

The wreath product is equipped with the quotient map $\alpha: A \wr G \rightarrow G$ defined by $\alpha(f\sigma) = \sigma$. The following lemmas describe two basic facts on embedding problems with wreath products.

Lemma 5.1. *Let $(\nu: \Gamma \rightarrow G, \alpha: A \wr G \rightarrow G)$ be a finite embedding problem for a profinite group Γ , assume $G \neq 1$, and let $\theta: \Gamma \rightarrow A \wr G$ be a weak solution. Then the only subgroup of A^1 that is normal in $\theta(\Gamma)$ is the trivial subgroup.*

Proof. For each $\sigma \in G$ choose $\gamma \in \Gamma$ such that $\nu(\gamma) = \sigma$, and let $f_\sigma = \theta(\gamma)\sigma^{-1}$, i.e. $\theta(\gamma) = f_\sigma\sigma$.

Assume that $B \leq A^1$ is normal in $\theta(\Gamma)$. Let $1 \neq \sigma \in G$. Then since

$$B = B^{f_\sigma\sigma} \leq (A^1)^{f_\sigma\sigma} = A^\sigma$$

we have

$$B \leq A^1 \cap A^\sigma = 1,$$

and hence $B = 1$. □

Lemma 5.2. *Let $K_0 \subseteq L$ be a finite Galois extension with Galois group $G = \text{Gal}(L/K_0)$ and let $\mu: \text{Gal}(K_0) \rightarrow G$ be the restriction map. Let A be a finite group that is regular over K_0 . Then the embedding problem*

$$(\mu: \text{Gal}(K_0) \rightarrow G, \alpha: A \wr G \rightarrow G)$$

is rational.

Proof. Since A is regular over K_0 there exists an absolutely irreducible polynomial $f(T, X) \in K_0[T, X]$ that is Galois over $K(T)$ and $\text{Gal}(f(T, X), K(T)) \cong A$.

Choose a basis $c_1, \dots, c_n$ of L/K_0 and let $\mathbf{t} = (t_1, \dots, t_n)$ be an n -tuple of variables. By [13], Lemma 3.1 (with $L_0 = L$ and $G_0 = 1$) there exist a field $\hat{F}$ such that

- (a) $\hat{F}$ is regular over L_0 ,
- (b) $\text{Gal}(\hat{F}/K_0(\mathbf{t})) \cong A \wr G$ and under this identification $\alpha: A \wr G \rightarrow G$ coincides with the restriction map $\text{Gal}(\hat{F}/K_0(\mathbf{t})) \rightarrow \text{Gal}(L_0/K_0)$.

In particular we get that (μ, α) is rational, as claimed. □

Remark 5.3. One can formulate and prove a much more general result than the above lemma. This generalization considers a split embedding problem $A \rtimes G_0 \rightarrow G_0$

instead of A , and gives a rational embedding problem with the twisted wreath product $A \wr_{G_0} G$ instead of $A \wr G$.

The proof of this generalization is a bit more technical, but still it uses only [13], Lemma 3.1. We will not use the generalization here, so we decided to omit it. For the full version see [2].


5.2. Proof of Theorem 1. Let K/K_0 be a proper separable PAC extension. We need to prove that the Galois closure of K/K_0 is K_{0s} .

We break the proof into two steps.

Step A. *If K/K_0 is a proper Galois PAC extension, then $K = K_{0s}$.*

Let $K_0 \subsetneq L \subseteq K$ be a finite Galois extension with a Galois group $G = \text{Gal}(L/K_0)$. Let N/K be a finite Galois extension with Galois group $B = \text{Gal}(N/K)$. It suffices to show that $B = 1$.

Identify B with a subgroup of $A = S_n$, for some sufficiently large n . Let $\nu: \text{Gal}(K_0) \rightarrow G$ and $\theta: \text{Gal}(K) \rightarrow B \leq A$ be the restriction maps. Since A is regular over any field [18], p. 272, Example 4, and in particular over K_0 , we get that the embedding problem (μ, α) is rational (Lemma 5.2). Hence, by definition, the double embedding problem



is rational. By the lifting property (Proposition 4.6) we can extend the weak solution θ of the lower embedding problem to a weak solution (θ, θ_0) of the double embedding problem. Now since $\text{Gal}(K) \triangleleft \text{Gal}(K_0)$ we have $B = \theta_0(\text{Gal}(K)) \triangleleft \theta_0(\text{Gal}(K_0))$. Thus $B = 1$ (Lemma 5.1), as needed.

Step B. *The general case.* Let M be the Galois closure of K/K_0 . We need to show that $M = K_{0s}$. By Theorem 6 (which was proved in [22] and will be reproved below) there exists M_0/K_0 that is linearly disjoint from K over K_0 and such that $M = KM_0$.

$$\begin{array}{ccc} K & \text{---} & M \\ | & & | \\ K_0 & \text{---} & M_0 \end{array}$$

In particular M/M_0 is a proper Galois extension. By [15], Lemma 2.1, M/M_0 is PAC. We get from the first step that $M = K_{0s}$. $\square$

5.3. PAC fields being non-PAC over any proper subfield. The following observation follows directly from Theorem 1.

Lemma 5.4. *Let $N \neq \tilde{\mathbb{Q}}$ be a Galois extension of $\mathbb{Q}$. Then N is a PAC extension of no proper subfield.*

Proof. If K is a proper subfield of N , then $\mathbb{Q} \subseteq K$. In particular N/K is Galois, and hence Theorem 1 implies that N/K is not PAC. $\square$

Some Galois extension of $\mathbb{Q}$ are known to be PAC as fields. Hence we get examples of PAC fields which are PAC extensions of no proper subfield.

Example i: The Galois hull $\tilde{\mathbb{Q}}[\sigma]$ of $\mathbb{Q}$ in $\tilde{\mathbb{Q}}(\sigma)$, for almost all $\sigma \in \text{Gal}(\mathbb{Q})^e$ [11], Theorem 18.10.2.

Example ii: $\mathbb{Q}_{\text{tr}}(i)$, where $\mathbb{Q}_{\text{tr}}$ is the maximal real Galois extension of $\mathbb{Q}$ and $i^2 = -1$ [20].

Example iii: The compositum $\mathbb{Q}_{\text{sym}}$ of all Galois extensions of $\mathbb{Q}$ with a symmetric Galois group [11], Theorem 18.10.3.

Over a finite field any infinite algebraic extension is PAC [11], Corollary 11.2.4. Thus we get

Example iv: Let N be an infinite extension of a finite field $\mathbb{F}_p$ which is not algebraically closed. Then N is a PAC field. However N is Galois over any subfield (since $\text{Gal}(\mathbb{F}_p) = \hat{\mathbb{Z}}$ is abelian). Hence, by Theorem 1, N is a PAC extension of no proper subfield.

5.4. Finite PAC Extensions – Proof of Theorem 3. Let K/K_0 be a finite extension. We need to prove that K/K_0 is PAC if and only if either K_0 is real closed and K is its algebraic closure or K_0 is a PAC field and K/K_0 is a finite purely inseparable extension.

Since an algebraically closed field is PAC over any infinite subset of it we have that indeed K_0 is real closed and K is its algebraic closure implies that K/K_0 is PAC. Moreover $[K : K_0] = 2$ (Artin-Schreier Theorem [18], VI§Corollary 9.3).

Let K_0 be PAC and K/K_0 a finite purely inseparable extension. Then [15], Corollary 2.3, asserts that K/K_0 is PAC.

For the other direction, assume that K/K_0 is a finite PAC extension. Let K_1 be the maximal separable extension of K_0 contained in K . Then K/K_1 is purely inseparable [18], V§6 Proposition 6.6. By [15], Corollary 2.3, K_1/K_0 is PAC, and in particular K_1 is a PAC field. If $K_1 = K_0$, we are done, since K/K_0 is then purely inseparable.

Assume $K_1 \neq K_0$. By Theorem 1, the Galois closure N of K_1/K_0 is the separable closure. Hence, by Artin-Schreier Theorem [18], VI§Corollary 9.3, N is, in fact, algebraically closed and K_0 is real closed (recall that $1 < [N : K_0] < \infty$). In particular, the characteristic of K is 0, and hence $K_1 = K$. $\square$

6. DESCENT FEATURES

6.1. Proof of Theorem 7. Let K/K_0 be a PAC extension and L/K a finite Galois extension. Assume $\text{Gal}(L/K) \leq G_0$, where G_0 is regular over K_0 .

We need to find a Galois extension L_0/K_0 such that $\text{Gal}(L_0/K_0) \leq G_0$ and $L = L_0K$.

$$\begin{array}{ccccc} & & K & \xrightarrow{G} & L \\ & & \downarrow & & \downarrow \\ K_0 & \xrightarrow{\quad} & L_0 \cap K & \xrightarrow{G} & L_0 \end{array}$$

The restriction map $\theta: \text{Gal}(K) \rightarrow G$ is a solution of the lower embedding problem of the rational double embedding problem

$$(10) \quad \begin{array}{ccccc} & & \text{Gal}(K) & & \\ & \swarrow \theta & \downarrow & \searrow & \\ & G & \text{Gal}(K_0) & & 1 \\ & \swarrow \theta_0 & \downarrow & \searrow & \\ G & \xrightarrow{\quad} & G_0 & \xrightarrow{\quad} & 1 \end{array}$$

It extends to a geometric weak solution (θ, θ_0) of (10) by the lifting property (Proposition 4.6). Let L_0 be the fixed field of $\ker(\theta_0)$. Then $\text{Gal}(L_0/K_0) = \theta_0(\text{Gal}(K_0)) \leq \text{Gal}(L_0/K_0)$. Since $\theta_0(\text{Gal}(K)) = \theta(\text{Gal}(K)) = G$ we get that $L = L_0K$. $\square$

6.2. Corollaries of Theorem 7. If the group G in Theorem 7 is regular over K_0 we can take $G = G_0$:

Corollary 6.1. *Let K/K_0 be a PAC extension. Let G be a finite Galois group over K that is regular over K_0 . Then G occurs as a Galois group over K_0 .*

Since every abelian group is regular over any field (see e.g. [11]), Proposition 16.3.5, we get the following

Corollary 6.2. *Let K/K_0 be a PAC extension. Then $K^{\text{ab}} = K_0^{\text{ab}}K$.*

From the fact that the symmetric group is regular over any field Theorem 7 gives a new proof of Theorem 6. This new proof provides an insight into Razon's original technical proof of Theorem 6 in [22].

Proof of Theorem 6. Let K/K_0 be a PAC extension and let L/K be a separable extension. We need to find a separable L_0/K_0 that is linearly disjoint from K over K_0 such that $L = L_0K$.

First assume that $[L : K]$ is finite. Let M be the Galois closure of L/K , $G = \text{Gal}(M/K)$, $G' = \text{Gal}(M/L)$. The action of G on the cosets $\Sigma = G/G'$ admits an embedding $i: G \rightarrow S_\Sigma$.

As S_Σ is regular over K_0 ([18], p. 272, Example 4), Theorem 7 gives a Galois extension M_0/K_0 with Galois group $H = \text{Gal}(M_0/K_0)$ such that $H \leq S_\Sigma$ and $G \leq H$ (since $M = M_0K$). Then H is transitive, since G is. Thus $(H : H') = |\Sigma| = [L : K]$, where H' is the stabilizer in H of the coset $G' \in \Sigma$. Also, as the subgroup $G' \leq G$ is the stabilizer in G of the coset $G' \in \Sigma$, it follows that $H' \cap G = G'$.

Let $L_0 \subseteq M_0$ be the corresponding fixed field of H' (i.e. $\text{Gal}(M_0/L_0) = H'$). So by the Galois correspondence $\text{Gal}(L) = \text{Gal}(L_0) \cap \text{Gal}(K) = \text{Gal}(L_0K)$, hence $L = L_0K$. In addition, L_0 is linearly disjoint from K , since $[L_0 : K_0] = (H : H') = [L : K]$, as needed.

The case where L/K is an infinite extension follows from Zorn's Lemma. The main point is that for a tower of algebraic extensions $L_1 \subseteq L_2 \subseteq L_3$, L_3/L_1 is separable if and only if both L_2/L_1 and L_3/L_2 are. The details can be found in [22]. $\square$

Remark 6.3. In the last proof H' was the stabilizer of a point of a subgroup of S_n . This stabilizer is, in general, not normal even if L/K is Galois. That is to say, L_0/K_0 need not be Galois, even if L/K is.

7. FIELDS WHICH ARE FINITE SEPARABLE/GALOIS EXTENSIONS OF NO PROPER SUBFIELD

In this section we prove the generalizations of Theorems 2 and 4. In particular we settle Problem 18.7.8 of [11] for finitely generated infinite fields. Before doing this we need a technical preparation about Hilbertian fields over subsets and their relation with PAC extensions.

7.1. Hilbertian Fields over Subsets. Let us start by introducing some notation. Let

$$f_1(T_1, \dots, T_e, X_1, \dots, X_n), \dots, f_m(T_1, \dots, T_e, X_1, \dots, X_n) \in K[\mathbf{T}, \mathbf{X}]$$

be irreducible polynomials and $g(\mathbf{T}) \in K[\mathbf{T}]$ nonzero. The corresponding **Hilbert set** is the set of all irreducible specializations $\mathbf{T} \mapsto \mathbf{a} \in K^e$ for $f_1, \dots, f_m$ under

which g does not vanish, i.e.

$$H_K(f_1, \dots, f_m; g) = \{\mathbf{a} \in K^r \mid \forall i \ f_i(\mathbf{a}, \mathbf{X}) \text{ is irreducible in } K[\mathbf{X}] \text{ and } g(\mathbf{a}) \neq 0\}.$$

Now K is **Hilbertian** if any Hilbert set is nonempty provided that $n = 1$ and $f_i = f_i(\mathbf{T}, X)$ is separable in X for each i . (Some authors use the terminology ‘ K is separably Hilbertian’.) A stronger property is that any Hilbert set for K is nonempty. We call such a field **s-Hilbertian**.

In case the characteristic of K is zero, these two properties coincide. If the characteristic of K is positive, there is a simple criterion for a Hilbertian field to be s-Hilbertian.

Theorem 7.1 (Uchida [11], Proposition 12.4.3). *Let K be a Hilbertian field of characteristic $p > 0$. Then K is s-Hilbertian if and only if K is imperfect.*

(Recall that K is imperfect if $[K : K^p] > 1$.)

Definition 7.2. A field E is said to be **Hilbertian over** a subset K if

$$H_E(f_1, \dots, f_m; g) \cap K^r \neq \emptyset$$

for any irreducible $f_1, \dots, f_m \in E[\mathbf{T}, X]$ that are separable in X and any nonzero $g(\mathbf{T}) \in E[\mathbf{T}]$. If furthermore $H_E(f_1, \dots, f_m; g) \cap K^r \neq \emptyset$ for any irreducible $f_1, \dots, f_m \in E[\mathbf{T}, \mathbf{X}]$, then we say that E is **s-Hilbertian over** K .

Note that a field K is Hilbertian (resp. s-Hilbertian) if and only if it is Hilbertian (resp. s-Hilbertian) over itself.

Jarden and Razon prove that if R is a ring with quotient field K and K is a countable Hilbertian field over R , then $K_s(\sigma)/R$ and $\tilde{K}(\sigma)$ are PAC for almost all $\sigma \in \text{Gal}(K)^e$ [15], Proposition 3.1.

A crucial observation for our applications is that the proof of [15], Proposition 3.1, gives the following stronger statement.

Theorem 7.3. *Let E be a countable field that is Hilbertian over a subset K . Then for almost all $\sigma \in \text{Gal}(E)^e$ the fields $E_s(\sigma)$ and $\tilde{E}(\sigma)$ are PAC over K .*

We shall use the last result to find new PAC extensions, and we start by finding Hilbertian fields over other fields.

Lemma 7.4. *Let K be an s-Hilbertian field over a subset S and E/K a purely transcendental extension. Then E is s-Hilbertian over S .*

Proof. Let $f_1(\mathbf{T}, \mathbf{X}), \dots, f_r(\mathbf{T}, \mathbf{X}) \in E[\mathbf{T}, \mathbf{X}]$ be irreducible polynomials and $0 \neq g(\mathbf{T}) \in E[\mathbf{T}]$. Since $E = K(u_\alpha \mid \alpha \in A)$, where $\{u_\alpha \mid \alpha \in A\}$ is a set of variables, we can assume that $f_i(\mathbf{T}, \mathbf{X}) = g_i(\mathbf{u}, \mathbf{T}, \mathbf{X})$, where

$$g_1(\mathbf{u}, \mathbf{T}, \mathbf{X}), \dots, g_r(\mathbf{u}, \mathbf{T}, \mathbf{X}) \in K[\mathbf{u}, \mathbf{T}, \mathbf{X}]$$

for some finite tuple of variables $\mathbf{u}$.

Since K is s -Hilbertian over S , there exists a tuple $\mathbf{a}$ of elements in S such that all $f_i(\mathbf{a}, \mathbf{X}) = g_i(\mathbf{u}, \mathbf{a}, \mathbf{X})$ are irreducible in $K[\mathbf{u}, \mathbf{X}]$ and $g(\mathbf{a}) \neq 0$. But the elements in $\{u_\alpha \mid \alpha \in A\}$ are algebraically independent, so all $f_i(\mathbf{a}, \mathbf{X}) = g_i(\mathbf{u}, \mathbf{a}, \mathbf{X})$ are irreducible in the larger ring $E[\mathbf{X}]$. $\square$

Proposition 7.5. *Let K be an s -Hilbertian field over a subset S and let E/K be a finitely generated extension. Then E is Hilbertian over S . Moreover, if E/K is also separable, then E is even s -Hilbertian over S .*

Proof. Choose a transcendence basis $\mathbf{t}$ for E/K , i.e., $K(\mathbf{t})/K$ is purely transcendental and $E/K(\mathbf{t})$ is finite. Let $H \subseteq E^r$ be a separable Hilbert set for E . By [11], Proposition 12.3.3, there exists a separable Hilbert set $H_1 \subseteq K(\mathbf{t})^r$ such that $H_1 \subseteq H$. By Lemma 7.4, we get that $H_1 \cap S^r \neq \emptyset$, and hence the assertion.

If E/K is also separable, then we can choose $\mathbf{t}$ to be a separating transcendence basis, that is, we can assume that $E/K(\mathbf{t})$ is separable. Now the same argument as above work for any Hilbert set $H \subseteq E^r$ (using [11], Corollary 12.2.3, instead of Proposition 12.3.3). $\square$

Combining the results that we attained so far, we enlarge the family of PAC extensions:

Theorem 7.6. *Let $e \geq 1$ be an integer, let K be a countable field which is s -Hilbertian over some subset S , and let E/K be a finitely generated extension. Then for almost all $\sigma \in \text{Gal}(E)^e$ the fields $E_s(\sigma)$ and $\tilde{E}(\sigma)$ are PAC over S .*

In particular, the result is valid when K is a countable s -Hilbertian field (and $S = K$).

Corollary 7.7. *Let $e \geq 1$ be an integer and let K be a finitely generated infinite field (over its prime field). Then for almost all $\sigma \in \text{Gal}(K)^e$ the field $K_s(\sigma)$ is a PAC extension of any subfield which is not algebraic over a finite field. Moreover, if K is of characteristic 0, then $K_s(\sigma)$ is also PAC over any subring.*

Proof. First assume that K is of characteristic 0. Then any ring contains $\mathbb{Z}$, so it suffices to show that $K_s(\sigma)/\mathbb{Z}$ is PAC for almost all $\sigma \in \text{Gal}(K)^e$. And indeed,

since $\mathbb{Q}$ is Hilbertian over $\mathbb{Z}$ and K is finitely generated over $\mathbb{Q}$, Theorem 7.6 implies that $K_s(\sigma)/\mathbb{Z}$ is PAC for almost all σ .

Next assume that the characteristic of K is $p > 0$. Since any field F which is not algebraic over $\mathbb{F}_p$ contains a rational function field $\mathbb{F}_p(t)$, it suffices to show that $K_s(\sigma)/\mathbb{F}_p(t)$ is PAC for almost all $\sigma \in \text{Gal}(K)^e$ and any $t \in K_s(\sigma) \setminus \tilde{\mathbb{F}}_p$.

Set $G = \text{Gal}(K)^e$ and let μ be its normalized Haar measure. For any $t \in K_s \setminus \tilde{\mathbb{F}}_p$ we define a subset $\Sigma_t \subseteq G$ as follows:

$$(11) \quad \Sigma_t = \{\sigma \in G \mid \text{if } t \in K_s(\sigma), \text{ then } K_s(\sigma)/\mathbb{F}_p(t) \text{ is PAC}\}.$$

We claim that $\mu(\Sigma_t) = 1$. Indeed, let $E = K(t)$. Then E/K is a finite separable extension. Let $H = \text{Gal}(E)^e$ be the corresponding open subgroup of G .

Note that $t \in K_s(\sigma)$ if and only if $\sigma \in H$. Then the definition of Σ_t implies that

$$\Sigma_t = (H \cap \Sigma_t) \cup (G \setminus H).$$

Hence it suffices to show that $\mu(H \cap \Sigma_t) = \mu(H)$, or equivalently, $\nu(H \cap \Sigma_t) = 1$, where ν denotes the normalized Haar measure on H .

Since $\mathbb{F}_p(t)$ is Hilbertian ([11], Theorem 13.3.5) and imperfect ([11], Lemma 2.7.2), Uchida's theorem implies that $\mathbb{F}_p(t)$ is s-Hilbertian. Also $E/\mathbb{F}_p(t)$ is finitely generated because K is.

Finally, since $H \cap \Sigma_t$ is the set of all $\sigma \in \text{Gal}(E)^e$ for which $E_s(\sigma)/\mathbb{F}_p(t)$ is PAC, and since $E_s = K_s$, Theorem 7.6 implies that $\nu(H \cap \Sigma_t) = 1$, as desired. $\square$

7.2. The Bottom Theorem. Now we are ready to address Problem 18.7.8 of [11], the so-called ‘bottom theorem’. Let K be a Hilbertian field and $e \geq 1$ an integer. The problem asks whether for almost all $\sigma = (\sigma_1, \dots, \sigma_e) \in \text{Gal}(K)^e$ the field $M = K_s(\sigma)$ has no cofinite proper subfield (that is, $N \subsetneq M$ implies $[M : N] = \infty$). Here the phrase ‘for almost all’ refers to the Haar measure on the profinite, and hence compact, group $\text{Gal}(K)$.

Note that the Hilbertian field $K = \mathbb{F}_p(t)$ has imperfect degree p , i.e., $[K : K^p] = p$. Moreover, the imperfect degree is preserved under separable extensions (see [11], Lemma 2.7.3), and hence every separable extension M/K satisfies $[M : M^p] = p$. In particular, M^p is a cofinite proper subfield of $M = K_s(\sigma)$ for all $\sigma \in \text{Gal}(K)^e$. Consequently, the problem requires a small modification, namely a separability assumption:

Conjecture 7.8. Let K be a Hilbertian field and $e \geq 1$ an integer. Then for almost all $\sigma \in \text{Gal}(K)^e$ the field $K_s(\sigma)$ is a finite separable extension of no proper subfield.

In [12] Haran proves an earlier version of this conjecture, namely with the additional assumption that $K \subseteq N$ (see also [11], Theorem 18.7.7).

We settle Conjecture 7.8 in the case K is a finitely generated infinite field (which is Hilbertian).

Theorem 7.9. *Conjecture 7.8 is true for a finitely generated infinite field K .*

Proof. Let $e \geq 1$ be an integer. Corollary 7.7 gives for almost all $\sigma \in \text{Gal}(K)^e$ that the field $K_s(\sigma)$ is a PAC extension of any subfield of it which is not algebraic over a finite field. We can also assume that $K_s(\sigma) \neq K_s$.

Therefore, if $N \subsetneq K_s(\sigma)$ is finite, then Theorem 3 implies that $K_s(\sigma)/N$ is purely inseparable. $\square$

7.3. Fields with no Galois Subfields. We strengthen Chatzidakis's result, Theorem 2, for finitely generated infinite fields.

Theorem 7.10. *Let K be a finitely generated infinite field and $e \geq 1$ an integer. Then for almost all $\sigma \in \text{Gal}(K)^e$ the field $K_s(\sigma)$ is a Galois extension of no proper subfield.*

Proof. Corollary 7.7 gives for almost all $\sigma \in \text{Gal}(K)^e$ that the field $K_s(\sigma)$ is a PAC extension of any subfield of it which is not algebraic over a finite field. We can also assume that $K_s(\sigma) \neq K_s$.

Therefore, Theorem 1 implies that $K_s(\sigma)$ is a Galois extension of no proper subfield. $\square$

REFERENCES

1. L. Bary-Soroker, *Projective Pairs of Profinite Groups*, Journal of Group Theory, in press.
2. L. Bary-Soroker, *Pseudo Algebraically Closed Extensions of Fields*, PhD thesis, Tel Aviv University (2008).
3. L. Bary-Soroker, *Dirichlet's Theorem for Polynomial Rings*, Proc. Amer. Math. Soc. **137**, 73–83 (2009).
4. L. Bary-Soroker, *On the Characterization of Hilbertian Fields*, Int. Math. Res. Not., Vol. 2008: article ID rnn089, 10 pages (2008).
5. L. Bary-Soroker and M. Jarden, *PAC Fields over Finitely Generated Fields*, Math. Z. **260** (2), 329–334 (2008).
6. L. Bary-Soroker and D. Kelmer, *On PAC extensions and scaled Trace forms*, Israel J. Math.
7. P. Dèbes, *Galois covers with prescribed fibers: the Beckmann-Black problem*, Ann. Scuola Norm. Sup. Pisa, Cl. Sci. **28** (1999), no. 4, 273–286.
8. A.J. Engler, *Fields with two incomparable Henselian valuation rings*, manuscripta mathematica **23** (1978), 373–385.
9. G. Faltings, *Endlichkeitssätze für abelsche Varietäten über Zahlkörpern*, Inv. Math. **73** (1983), no. 3, 349–366.
10. M. D. Fried and M. Jarden, *Field Arithmetic*, Ergebnisse der Mathematik (3) **11**, Springer-Verlag, Heidelberg, 1986.
11. M. D. Fried and M. Jarden, *Field Arithmetic*, second ed., revised and enlarged by M. Jarden, Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics, vol. 11, Springer-Verlag, Berlin, 2005.
12. D. Haran, *The Bottom Theorem*, Arch. Math. (Basel) **45** (1985), no. 3, 229–231.
13. D. Haran, *Hilbertian Fields under Separable Algebraic Extensions*, Inv. Math. **137** (1999), 85–112.
14. M. Jarden, *PAC Fields over Number Fields*, Journal de Theorie des Nombres de Bordeaux **18** (2006), no. 2, 371–377.

15. M. Jarden and A. Razon, *Pseudo Algebraically Closed Fields over Rings*, Israel J. Math. **86** (1994), no. 1-3, 25–59.
16. M. Jarden and A. Razon, *Rumely's Local Global Principle for Algebraic PSC Fields over Rings*, Trans. Amer. Math. Soc. **350** (1998), no. 1, 55–85.
17. M. Jarden and A. Razon, *Rumely Local Global Principle for Weakly PSC Fields over Holomorphic Domains*, manuscript, 2006.
18. S. Lang, *Algebra*, Revised third edition, Graduate Texts in Mathematics, **211**, Springer-Verlag, New York, 2002.
19. Y. V. Matiyasevich, *Hilbert's tenth problem*, Translated from the 1993 Russian original by the author. With a foreword by Martin Davis. Foundations of Computing Series. MIT Press, Cambridge, MA, 1993.
20. F. Pop, *Embedding Problems over Large Fields*, Ann. of Math. (2) **144** (1996), no. 1, 1–34.
21. A. Razon, *Abundance of Hilbertian Domains*, Manuscripta Mathematica **94** (1997), 531–542.
22. A. Razon, *Splittings of $\tilde{\mathbb{Q}}/\mathbb{Q}$* , Arch. Math. (Basel) **74** (2000), no. 4, 263–265.
23. R. Rumely, *Arithmetic over the ring of all algebraic integers*, J. Reine Angew. Math. **368** (1986), 127–133.
24. P. Samuel, *Lectures on Old and New Results on Algebraic Curves*, Tata Institute of Fundamental Research, Bombay, 1966.

THE RAYMOND AND BEVERLY SACKLER SCHOOL OF MATHEMATICAL SCIENCES, TEL-AVIV UNIVERSITY
 E-mail address: barylior@post.tau.ac.il